

Effects of Sea Breeze on City Ventilation - Important for Air Ventilation Assessments?



11
102
1004

Leibniz
Universität
Hannover

Tobias Gronemeier¹, Siegfried Raasch¹, Edward Ng²
¹Leibniz University Hannover, Germany ²Chinese University of Hong Kong, China

Why we did it

- As urban population increases, preserving an acceptable city climate becomes a major challenge of future city planning
 - Air Ventilation Assessments (AVAs) help city planners to predict city ventilation of planned construction sites
 - To sufficiently predict ventilation, an AVA must cover the real atmospheric situation
 - Current AVA focus only on neutral conditions
- Is this sufficient for summery weak-wind conditions in a coastal city area?

How we did it

- LES simulation of Kowloon peninsula (Hong Kong), using the model PALM
- Summer weak-wind condition: easterly background wind: 1.5 ms⁻¹, fixed surface heat flux: 200 Wm⁻²
- Two cases:
 - homogeneous heating throughout domain
 - sea-breeze case where only land is heated
- Divide city into 3 regions (according to [2]): C1: SW ventilated; C2: weakly ventilated; C3: SE ventilated
- Passive scalar released at surface within city area




Figure 1: Model domain with terrain height in m and region definition.

How is the ventilation?

- Sea breeze penetrates city area along the entire coastline, forming a convergence zone above the city
- Higher V_r at west coast due to lower building density and flat terrain
- Comparison shows higher V_r in sea-breeze case especially in western part of Kowloon

Definitions:

$$V_r = \frac{V_{2m}}{V_{ref}}$$

$$V_{ref} = V_{2m} \text{ (outside of city area) [1]}$$

$$V_{r, norm} = \frac{V_r \text{ (sea-breeze case)}}{V_r \text{ (homog. heating case)}}$$




Figure 2: Velocity ratio V_r and wind vectors of the sea-breeze case (left), and normalized velocity ratio $V_{r, norm}$ (right).

Is the pollution dispersion influenced by sea-breeze?

- Scalar concentration differs significantly between cases
- Strong west-east gradient in sea-breeze case, north-south gradient in homog. heating case
- Different wind fields yield large differences in s^* between cases
- Depending on case, high s^* values are observed in different city areas

Definitions:

$$s^* = \frac{s}{Q_s t_e}$$

t_e : emission time
 Q_s : emission rate




Figure 3: Normalized scalar concentration s^* and wind vectors for the (left) sea-breeze case and (right) homogeneous-heating case.

Looking at different city regions

- C1, C2, C3 correspond to different city regions according to [2]
 - Mean wind direction (dir, Tab.1) agrees with classification made by [2] for sea-breeze case but not for homog. heating case
 - Although C1 has highest ventilation, pollution is also highest
- Between Kowloon and Hong Kong Island, complex wind circulation transports pollution over sea where it re-enters the city area
- Vanishingly low correlation between V_r and mean building height H_{avg} confirm that H_{avg} plays a minor role for city ventilation (Fig.4, see also [1])

	V_r	s^*	dir
C1	1.74 : 1.46	0.0023 : 0.0023	224° (SW) : 149° (SE)
C2	1.29 : 1.21	0.0022 : 0.0022	165° (S) : 168° (S)
C3	1.38 : 1.34	0.0020 : 0.0020	137° (SE) : 145° (SE)

Table 1: Mean values within regions C1-C3; left: sea-breeze case, right: homog. heating case.




Figure 4: Correlation between velocity ratio V_r and (left) mean building height H_{avg} and (right) plan-area index PAI; each data point represents a 100m x 100m area within the specified region; sea-breeze case.

Let's summarize

- Ventilation changes significantly between sea-breeze case and homogeneously heated case in strength and direction
 - More complex wind circulation lead to differences in pollution concentration (W-E gradient instead of N-S)
 - Main wind direction from measurements can only be reproduced if sea-breeze is considered
- It is essential to cover sea-breeze effects if a sufficient analysis of the city ventilation is focused during summery weak-wind conditions!

What's next

- Further detailed analysis of wind system between Kowloon and Hong Kong Island should reveal more details of sea-breeze effects on ventilation
- Compare results with real-world measurements in Hong Kong
- Using PALM's new nesting methods, a larger area can be simulated to study effects of large-scale wind systems (see also poster 1D-51)

References & Acknowledgments

- [1] Gronemeier, T., S. Raasch, E. Ng (2017). Effects of unstable stratification on ventilation in Hong Kong. Atmosphere, 8, 1-15, doi:10.3390/atmos8090168
- [2] Ng, E., V. Cheng, C. Chan (2008). Urban climatic map and standards for wind environment - feasibility study. Technical Input Report No. 1: Methodologies and findings of user's wind comfort level survey. Hong Kong Planning Department
- This study was supported by a research grant (14408214) from General Research Fund of Hong Kong Research Grants Council (HK RGC-GRF). All simulations were carried out on the Cray CX-40 systems of the North-German Supercomputing Alliance (HLRN). T. Gronemeier was supported by MOSAIK, funded by the German Federal Ministry of Education and Research (BMBF) under grant 01LP1601A (<http://www.fona.de>)

